



INVESTOR IN PEOPLE


The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed 
Dated 10 June 2003



17SEP02 E748684-1/001298
P01/7700 0000-0221477.3

1477

The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

1. Your reference

PC25375

2. Patent application number

(The Patent Office will fill in this part)

16 SEP 2002

0221477.3

3. Full name, address and postcode of the or of each applicant (*underline all surnames*)

PFIZER LIMITED
Ramsgate Road,
Sandwich,
Kent, CT13 9NJ

689 2673001

Patents ADP number (*if you know it*)

United Kingdom

If the applicant is a corporate body, give the country/state of its incorporation

4. Title of the invention

CHEMICAL COMPOUNDS

5. Name of your agent (*if you have one*)

Mr. G.M. Lane

"Address for service" in the United Kingdom to which all correspondence should be sent (*including the postcode*)

UK Patent Department
Ramsgate Road,
Sandwich, Kent,
CT13 9NJ
United Kingdom

8426322001

Patents ADP number (*if you know it*)

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (*if you know it*) the or each application number

Country

Priority application number
(*if you know it*)

Date of filing
(*day / month / year*)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(*day / month / year*)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (*Answer 'Yes' if:*

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is not named as an applicant, or
 - c) any named applicant is a corporate body.
- See note (d))

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description	24
Claim(s)	3
Abstract	1
Drawing(s)	0

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination and search (*Patents Form 9/77*)

Request for substantive examination (*Patents Form 10/77*)

Any other documents
(please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature



Date
16 September 2002

12. Name and daytime telephone number of person to contact in the United Kingdom

Mr. G.M. Lane

01304.642817

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

CHEMICAL COMPOUNDS


This invention relates to pyrazole derivatives, to their use in medicine, to compositions containing them, to processes for their preparation and to intermediates used in such processes.

Reverse transcriptase is implicated in the infectious lifecycle of Human Immunodeficiency Virus (HIV). Compounds which interfere with the function of this enzyme have shown utility in the treatment of conditions caused by HIV and genetically related retroviruses, such as Acquired Immune Deficiency Syndrome (AIDS). There is a constant need to provide new and better modulators, especially inhibitors, of HIV reverse transcriptase, since the virus is able to mutate, becoming resistant to the effects of known modulators.

A class of N-phenylpyrazoles which act as reverse transcriptase inhibitors are disclosed in *J. Med. Chem.*, 2000, **43**, 1034. Antiviral activity is ascribed to a class of N-(hydroxyethyl)pyrazole derivatives in US patent number 3,303,200. International Application No. PCT/IB02/01234, unpublished at the filing date of the instant application, generically embraces, but does not specifically disclose, the compound of formula (I) below.

The compounds of the invention bind to the enzyme reverse transcriptase and are modulators, especially inhibitors, thereof.

According to the invention there is thus provided the compound of formula (I)



or a pharmaceutically acceptable salt, solvate or derivative thereof.

The pharmaceutically acceptable salts of the compounds of formula (I) include the acid addition and the base salts thereof.



Suitable acid addition salts are formed from acids which form non-toxic salts and examples are the hydrochloride, hydrobromide, hydroiodide, chloride, bromide, iodide, sulphate, bisulphate, nitrate, phosphate, hydrogen phosphate, acetate, fumarate, pamoate, aspartate, besylate, carbonate, bicarbonate/, camsylate, D and L-lactate, D and L-tartrate, esylate, mesylate, malonate, orotate, gluceptate, methylsulphate, stearate, glucuronate, 2-napsylate, tosylate, hibenazate, nicotinate, isethionate, malate, maleate, citrate, gluconate, succinate, saccharate, benzoate, esylate, and pamoate salts.

10

Suitable base salts are formed from bases which form non-toxic salts and examples are the sodium, potassium, aluminium, calcium, magnesium, zinc, choline, diolamine, olamine, arginine, glycine, tromethamine, benzathine, lysine, meglumine and diethylamine salts.

15

For reviews on suitable salts see Berge *et al*, J. Pharm. Sci., 66, 1-19, 1977 and Bighley *et al*, Encyclopedia of Pharmaceutical Technology, Marcel Dekker Inc, New York, 1996, Vol 13, pp453-497

20 The pharmaceutically acceptable solvates of the compounds of formula (I) include the hydrates thereof.

The compound of formula (I) may be modified to provide pharmaceutically acceptable derivatives thereof at any of the functional groups in the compound.

25 Examples of such derivatives are described in: Drugs of Today, Volume 19, Number 9, 1983, pp 499 – 538; Topics in Chemistry, Chapter 31, pp 306 – 316; and in "Design of Prodrugs" by H. Bundgaard, Elsevier, 1985, Chapter 1 (the disclosures in which documents are incorporated herein by reference) and include: esters, carbonate esters, hemi-esters, phosphate esters, nitro esters, sulfate
30 esters, sulfoxides, amides, sulphonamides, carbamates, azo-compounds, phosphamides, glycosides, ethers, acetals and ketals.

The invention encompasses all isomers of the compound of formula (I) and pharmaceutically acceptable salts, solvates or derivatives thereof, including all
35 geometric, tautomeric and optical forms, and mixtures thereof (e.g. racemic mixtures).

Separation of diastereoisomers may be achieved by conventional techniques, e.g. by fractional crystallisation, chromatography or high performance liquid



chromatography (HPLC) of a stereoisomeric mixture of compounds. An individual enantiomer of a compound may also be prepared from a corresponding optically pure intermediate or by resolution, such as by HPLC of the corresponding racemate using a suitable chiral support, or by fractional crystallisation of the diastereoisomeric salts formed by reaction of the corresponding racemate with a suitable optically active acid or base, as appropriate.

The compound of formula (I) and pharmaceutically acceptable salts, solvates or derivatives thereof may have the ability to crystallize in more than one form, a characteristic known as polymorphism, and all such polymorphic forms ("polymorphs") are encompassed within the scope of the invention. Polymorphism generally can occur as a response to changes in temperature or pressure or both, and can also result from variations in the crystallization process. Polymorphs can be distinguished by various physical characteristics, and typically the x-ray diffraction patterns, solubility behaviour, and melting point of the compound are used to distinguish polymorphs.

The compound of formula (I), pharmaceutically acceptable salts, solvates and derivatives thereof, isomers thereof, and polymorphs thereof, are hereinafter referred to as the compounds of the invention.

Preferred compounds of the invention are the compound of formula (I) and its pharmaceutically acceptable salts and solvates thereof.

The most preferred compound of the invention is the compound of formula (I).

The compounds of the invention exhibit advantageous properties, including excellent metabolic stability, leading to excellent pharmacokinetic properties. In addition, the compounds of the invention may have advantages over those of the prior art with regard to a number of other useful properties, such as potency, duration of action, spectrum of activity, side effect profile, solubility, chemical stability, and so on.


The compounds of the invention may be prepared by any method known in the art for the preparation of compounds of analogous structure. The compounds of the invention can be prepared by the procedures described in the methods below, or by the specific methods described in the Examples, or by similar methods to either. The invention also encompasses any one or more of these



processes for preparing the compounds of the invention, in addition to any novel intermediates used therein.

5 The compound of formula (I) may be prepared according to the route shown in Scheme 1 below.

Scheme 1




10 In Scheme 1, the compound of formula (I) may be prepared by condensation of
the compound of formula (II) with 2-hydroxyethylhydrazine of formula (V) or a salt
or hydrate thereof, optionally in the presence of an acid or a base, the base
preferably being a tertiary amine base such as triethylamine and the acid
preferably being acetic acid. In a typical procedure, a solution of the compound
15 of formula (II) in a suitable solvent, such as acetic acid, is treated with the
hydrazine of formula (V), or the salt or hydrate thereof, and, if used, the
appropriate acid or base, at a temperature of from room temperature to the reflux
temperature of the solvent. In a preferred procedure, the reaction is carried out
at room temperature.

20


Functional equivalents of the compound of formula (II) may also be used in this
reaction. These include compounds of formulae (VI) or (VII), in which L^1 and L^2 ,



respectively, are each suitable leaving groups, preferably $-N(C_1-C_6 \text{ alkyl})_2$, most preferably $-N(CH_3)_2$.



(VI)



(VII)

5


Thus, the compound of formula (I) may be prepared by condensation of a compound of formulae (VI) or (VII) with the compound of formula (V), or a salt or hydrate thereof, optionally in the presence of an acid or a base, the base preferably being a tertiary amine base such as triethylamine and the acid preferably being acetic acid. In a typical procedure, a solution of the compound of formula (VI) or (VII) in a suitable solvent, such as ethanol, is treated with the compound of formula (V), or the salt or hydrate thereof, and, if used, the appropriate acid or base, at a temperature of from room temperature to the reflux temperature of the solvent. In a preferred procedure, the reaction mixture is heated under reflux.

15


Compounds of formula (VI) in which L^1 is dimethylamino may be prepared by the reaction of the compound of formula (VIII) with an appropriately substituted formamide acetal at an elevated temperature, preferably at about 100°C .

Compounds of formula (VII) in which L^1 is dimethylamino may be prepared by the reaction of the compound of formula (IX) under the same conditions.

20




(VIII)



(IX)




The compound of formula (VIII) is either commercially available or may be prepared by the reaction of the compound of formula (X)



5


with the compound of formula (XI)



10 In a typical procedure, a solution of the compound of formula (XI) in a suitable solvent, such as acetone, is treated with a suitable base, such as caesium carbonate, and the compound of formula (X). In a preferred procedure, the reaction mixture is heated, for example under reflux. Optionally, a nucleophilic catalyst such as sodium iodide or tetrabutylammonium iodide may be added.

15

The compound of formula (IX) is either commercially available or may be prepared from the compound of formula (XII)



20

and the compound of formula (XI) in the same way that the compound of formula (VIII) may be prepared from the compound of formula (X).


25 The compound of formula (II) may be prepared by the reaction of the compound of formula (III) with the compound of formula (XI).

30 In a typical procedure, a solution of the compound of formula (III) in a suitable solvent such as acetone is treated with the compound of formula (XI) and a suitable base, such as potassium or caesium carbonate, and heated, preferably under reflux. Optionally, a nucleophilic catalyst such as sodium iodide or tetrabutylammonium iodide may be added.




The compound of formula (III) is either commercially available or may be prepared by reaction of the compound of formula (IV) with a chlorinating reagent. In a typical procedure, a cooled solution of the compound of formula (IV) in a suitable solvent such as acetonitrile is treated first with tetrabutylammonium bromide and chlorotrimethylsilane and then dry dimethylsulphoxide. In another typical procedure, the compound of formula (IV) is treated with sulphuryl chloride, optionally in the presence of a suitable solvent such as dichloromethane.

- 10 The compound of formula (I) may also be prepared by reaction of the pyrazole of formula (XIII)



with an alkylating agent of formula (XIV)

15



or a protected derivative thereof.

- 20 In a typical procedure, a solution of the pyrazole of formula (XIII) in a suitable solvent such as ethanol or N,N-dimethylformamide is treated with an alkylating agent of formula (XIV) such as a protected hydroxyethyl bromide and a base such as sodium ethoxide or sodium hydride and heated at a temperature of from 0°C to the reflux temperature of the solvent. A preferred combination is N,N-dimethylformamide as the solvent, sodium hydride as the base, 0°C as the temperature and 2-(2-bromoethoxy)tetrahydro-2H-pyran as the alkylating agent.
- 25

As will be appreciated by the skilled artisan, in the alkylation of the pyrazole of formula (XIII) it may be necessary or desirable to protect the OH group of the compound of formula (XIV), in which case the compound of formula (I) is finally prepared by deprotection of the corresponding compound bearing an -OP¹ group, wherein P¹ is a suitable protecting group. Examples of suitable protecting groups

30



will be apparent to the skilled person; see, for instance, 'Protecting groups in Organic Synthesis (Second Edition)' by Theodora W. Green and Peter G. M. Wuts, 1991, John Wiley and Sons (in particular pages 10 - 118, relating to protection for the hydroxyl group), incorporated herein by reference. Such
5 compounds bearing an -OP¹ group may be prepared using the routes described above, *mutatis mutandis*.

Compounds of formulae (IV) and (V) are either commercially available, known from the literature or easily prepared by methods well known to those skilled in
10 the art.

The compounds of the invention can be administered alone, but will generally be administered in admixture with a suitable pharmaceutical excipient, diluent or carrier selected with regard to the intended route of administration and standard
15 pharmaceutical practice.

For example, the compounds of the invention can be administered orally, buccally or sublingually in the form of tablets, capsules, multi-particulates, gels, films, ovules, elixirs, solutions or suspensions, which may contain flavouring or
20 colouring agents, for immediate-, delayed-, modified-, sustained-, pulsed- or controlled-release applications. The compounds of the invention may also be administered as fast-dispersing or fast-dissolving dosage forms or in the form of a high energy dispersion or as coated particles. Suitable formulations of the compounds of the invention may be in coated or uncoated form, as desired.

Such solid pharmaceutical compositions, for example, tablets, may contain excipients such as microcrystalline cellulose, lactose, sodium citrate, calcium carbonate, dibasic calcium phosphate, glycine and starch (preferably corn, potato or tapioca starch), disintegrants such as sodium starch glycollate, croscarmellose
30 sodium and certain complex silicates, and granulation binders such as polyvinylpyrrolidone, hydroxypropylmethylcellulose (HPMC), hydroxypropylcellulose (HPC), sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, stearic acid, glyceryl behenate and talc may be included.

35

General Example

A formulation of the tablet could typically contain from 0.01mg to 500mg of active compound whilst tablet fill weights may range from 50mg to 1000mg. An example of a formulation for a 10mg tablet is illustrated below:



	<u>Ingredient</u>	<u>%w/w</u>
	Compound of the invention	10.000*
	Lactose	64.125
5	Starch	21.375
	Croscarmellose sodium	3.000
	Magnesium Stearate	1.500

* Quantity adjusted in accordance with drug activity.

10

The tablets are manufactured by a standard process, for example, direct compression or a wet or dry granulation process. The tablet cores may be coated with appropriate overcoats.

- 15 Solid compositions of a similar type may also be employed as fillers in gelatin or HPMC capsules. Preferred excipients in this regard include lactose, starch, a cellulose, milk sugar or high molecular weight polyethylene glycols. For aqueous suspensions and/or elixirs, the compounds of the invention may be combined with various sweetening or flavouring agents, colouring matter or dyes, with
- 20 emulsifying and/or suspending agents and with diluents such as water, ethanol, propylene glycol and glycerin, and combinations thereof.

- The compounds of the invention can also be administered parenterally, for example, intravenously, intra-arterially, intraperitoneally, intrathecally,
- 25 intraventricularly, intraurethrally, intrasternally, intracranially, intramuscularly or subcutaneously, or they may be administered by infusion or needleless injection techniques. For such parenteral administration they are best used in the form of a sterile aqueous solution which may contain other substances, for example, enough salts or glucose to make the solution isotonic with blood. The aqueous
- 30 solutions should be suitably buffered (preferably to a pH of from 3 to 9), if necessary. The preparation of suitable parenteral formulations under sterile conditions is readily accomplished by standard pharmaceutical techniques well-known to those skilled in the art.

- 35 For oral and parenteral administration to human patients, the daily dosage level of the compounds of the invention will usually be from 0.01 to 30 mg/kg, preferably from 0.01 to 5 mg/kg (in single or divided doses).



Thus tablets or capsules of the compound of the invention may contain from 1 to 500 mg of active compound for administration singly or two or more at a time, as appropriate. The physician in any event will determine the actual dosage which will be most suitable for any individual patient and it will vary with the age, weight and response of the particular patient. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited and such are within the scope of this invention. The skilled person will appreciate that, in the treatment of certain conditions the compounds of the invention may be taken as a single dose as needed or desired.

The compounds of invention can also be administered intranasally or by inhalation and are conveniently delivered in the form of a dry powder inhaler or an aerosol spray presentation from a pressurised container, pump, spray, atomiser or nebuliser, with or without the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, a hydrofluoroalkane such as 1,1,1,2-tetrafluoroethane (HFA 134A [trade mark]) or 1,1,1,2,3,3,3-heptafluoropropane (HFA 227EA [trade mark]), carbon dioxide or other suitable gas. In the case of a pressurised aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. The pressurised container, pump, spray, atomiser or nebuliser may contain a solution or suspension of the active compound, e.g. using a mixture of ethanol and the propellant as the solvent, which may additionally contain a lubricant, e.g. sorbitan trioleate. Capsules and cartridges (made, for example, from gelatin) for use in an inhaler or insufflator may be formulated to contain a powder mix of a compound of the invention and a suitable powder base such as lactose or starch.

Alternatively, the compounds of the invention can be administered in the form of a suppository or pessary, or they may be applied topically in the form of a gel, hydrogel, lotion, solution, cream, ointment or dusting powder. The compounds of the invention may also be dermally or transdermally administered, for example, by the use of a skin patch. They may also be administered by the pulmonary or rectal routes.

They may also be administered by the ocular route. For ophthalmic use, the compounds can be formulated as micronised suspensions in isotonic, pH adjusted, sterile saline, or, preferably, as solutions in isotonic, pH adjusted, sterile saline, optionally in combination with a preservative such as a benzylalkonium chloride. Alternatively, they may be formulated in an ointment such as petrolatum.



For application topically to the skin, the compounds of the invention can be formulated as a suitable ointment containing the active compound suspended or dissolved in, for example, a mixture with one or more of the following: mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, they can be formulated as a suitable lotion or cream, suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, a polyethylene glycol, liquid paraffin, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

The compounds of the invention may also be used in combination with a cyclodextrin. Cyclodextrins are known to form inclusion and non-inclusion complexes with drug molecules. Formation of a drug-cyclodextrin complex may modify the solubility, dissolution rate, bioavailability and/or stability property of a drug molecule. Drug-cyclodextrin complexes are generally useful for most dosage forms and administration routes. As an alternative to direct complexation with the drug the cyclodextrin may be used as an auxiliary additive, e.g. as a carrier, diluent or solubiliser. Alpha-, beta- and gamma-cyclodextrins are most commonly used and suitable examples are described in WO-A-91/11172, WO-A-94/02518 and WO-A-98/55148.

It is to be appreciated that all references herein to treatment include curative, palliative and prophylactic treatment.

Oral administration is preferred.

Included within the scope of the invention are embodiments comprising the co-administration of a compound of the invention with one or more additional therapeutic agents, and compositions containing a compound of the invention along with one or more additional therapeutic agents. Such a combination therapy is especially useful for the prevention and/or treatment of infection by HIV and related retroviruses which may evolve rapidly into strains resistant to any monotherapy. Alternatively, additional therapeutic agents may be desirable to treat diseases and conditions which result from or accompany the disease being treated with the compound of the invention. For example, in the treatment of an HIV or related retroviral infection, it may be desirable to additionally treat opportunistic infections, neoplasms and other conditions which occur as a result of the immuno-compromised state of the patient being treated.



Preferred combinations of the invention include simultaneous or sequential treatment with a compound of the invention and one or more:

- 5 (a) reverse transcriptase inhibitors such as abacavir, adefovir, didanosine, lamivudine, stavudine, zalcitabine and zidovudine;
- (b) non-nucleoside reverse transcriptase inhibitors such as capavirine, delavirdine, efavirenz, and nevirapine;
- (c) HIV protease inhibitors such as indinivir, nelfinavir, ritonavir, and
- 10 saquinavir;
- (d) CCR5 antagonists such as TAK-779 or UK-427,857;
- (e) CXCR4 antagonists such as AMD-3100;
- (f) integrase inhibitors, such as L-870,810 or S-1360;
- (g) inhibitors of viral fusion such as T-20;
- 15 (h) investigational drugs such as trizivir, KNI-272, amprenavir, GW-33908, FTC, PMPA, MKC-442, MSC-204, MSH-372, DMP450, PNU-140690, ABT-378, KNI-764, DPC-083, TMC-120 or TMC-125;
- (i) antifungal agents, such as fluconazole, itraconazole or voriconazole; or
- (j) antibacterial agents, such as azithromycin.

20

The activity of the compounds of the invention as reverse transcriptase inhibitors may be measured using the following assay.

Inhibition of HIV-1 reverse transcriptase enzyme

- 25 Using purified recombinant HIV-1 reverse transcriptase (RT, EC, 2.7.7.49) obtained by expression in Escherichia Coli, a 96-well plate assay system is established for assaying a large number of samples using either the Poly(rA)-oligo(dT) Reverse Transcriptase [3H]-SPA enzyme assay system (Amersham NK9020) or the [3H]-flashplate enzyme assay system (NEN - SMP 103) and
- 30 following the manufacturer's recommendations. The compounds are dissolved in 100% DMSO and diluted with the appropriate buffer to a 5% final DMSO concentration. The inhibitory activity is expressed in percent inhibition relative to DMSO control. The concentration at which compound inhibits reverse transcriptase by 50% is expressed as the IC_{50} of the compound.

35

The compound of Example 1, when tested according to the above procedure, had an IC_{50} value of 295 nanomolar.



The metabolism of the compounds of the invention may be measured using the following assays.

A. Metabolism in human liver microsomes and Supermix™

- 5 The metabolic vulnerability of the compounds of the invention in microsomes and Supermix™ may be assayed as follows.

10 The microsomal fraction is isolated from several human livers and the P450 content determined. Supermix™ is obtained from Gentest. Human microsomes (0.5µM cytochrome P450) and Supermix™ (0.05µM cytochrome P450) are added to an incubation media containing 50mM phosphate buffer (pH7.4), 5mM MgCl₂, 1mM NADP and an NADPH generating system consisting of isocitrate and isocitrate dehydrogenase. The substrate concentration is 1µM and incubations are carried out at 37°C for 1 hour. Samples are taken at time points
15 throughout this period and analysed using hplc-ms-ms assay.

The compound of Example 1, when tested according to the above procedure, had a t_{1/2} value of >120 minutes (both human microsomal and Supermix™).

20 B. Metabolism in Human hepatocytes.

The metabolic vulnerability of the compounds of the invention in human hepatocytes may be assayed as follows.

Cryopreserved human hepatocytes are obtained from *In vitro* Technologies, Inc.
25 The hepatocytes are thawed and suspended at 1 million cells/ml in 50% Krebs-Heinsleit buffer : 50% Williams E media containing 10% foetal bovine serum. The substrate concentration is 3µM and incubations are carried out at 37°C for 3 hours. Samples are taken at time points throughout this period and analysed using hplc-ms-ms assay.

30

The compound of Example 1, when tested according to the above procedure, had an unbound hepatocyte clearance value of <9 ml/min/kg.



Thus the invention provides:

- (i) the compound of formula (I) or a pharmaceutically acceptable salt, solvate or derivative thereof;
- 5 (ii) a process for the preparation of the compound of formula (I) or a pharmaceutically acceptable salt, solvate or derivative thereof;
- (iii) a pharmaceutical composition including the compound of formula (I) or a pharmaceutically acceptable salt, solvate or derivative thereof, together with a pharmaceutically acceptable excipient, diluent or carrier;
- 10 (iv) the compound of formula (I) or a pharmaceutically acceptable salt, solvate or composition thereof, for use as a medicament;
- (v) the compound of formula (I) or a pharmaceutically acceptable salt, solvate or composition thereof, for use as a reverse transcriptase inhibitor or modulator;
- 15 (vi) the compound of formula (I) or a pharmaceutically acceptable salt, solvate or composition thereof, for use in the treatment of an HIV or genetically-related retroviral infection, or a resulting acquired immune deficiency syndrome (AIDS);
- 20 (vii) the use of the compound of formula (I) or of a pharmaceutically acceptable salt, solvate or composition thereof, for the manufacture of a medicament having reverse transcriptase inhibitory or modulating activity;
- (viii) the use of the compound of formula (I) or of a pharmaceutically acceptable salt, solvate or composition thereof, for the manufacture of a medicament for the treatment of an HIV or genetically-related retroviral infection, or a resulting acquired immune deficiency syndrome (AIDS);
- 25 (ix) a method of treating an HIV or a genetically-related retroviral infection, or a resulting acquired immune deficiency syndrome (AIDS), comprising administering an effective amount of the compound of formula (I) or a pharmaceutically acceptable salt, solvate or composition thereof; and
- 30 (xi) certain novel intermediates disclosed herein.




The following Examples illustrate the preparation of the compounds of formula (I). The synthesis of certain intermediates used therein are described in the Preparations section that follows the Examples.

- 5 ¹H Nuclear magnetic resonance (NMR) spectra were in all cases consistent with the proposed structures. Characteristic chemical shifts (δ) are given in parts-per-million (ppm) downfield from tetramethylsilane using conventional abbreviations for designation of major peaks, e.g.: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; br, broad. The following abbreviations have been used: HRMS,
- 10 high resolution mass spectrometry; hplc, high performance liquid chromatography; nOe, nuclear Overhauser effect; m.p., melting point; CDCl₃, deuteriochloroform; D₆-DMSO, deuterodimethylsulphoxide; CD₃OD, deuteromethanol. Where thin layer chromatography (TLC) has been used it refers to silica gel TLC using silica gel 60 F₂₅₄ plates, R_f being the distance
- 15 travelled by a compound divided by the distance travelled by the solvent front on the TLC plate.



EXAMPLE 1

5-([3-Cyclopropyl-1-(2-hydroxyethyl)-5-methyl-1H-pyrazol-4-yl]oxy)isophthalonitrile



- 5 2-Hydroxyethylhydrazine (1.15ml, 16.9mmol) was added to a solution of the diketone from Preparation 7 (4.1g, 15.4mmol) in acetic acid (40ml) under nitrogen at room temperature. After stirring for 18 hours, the mixture was concentrated under reduced pressure and the residual oil was purified by flash chromatography on silica gel eluting with ethyl acetate:pentane (50:50 changing to 100:0, by volume) to provide samples of the two regioisomers which required further purification.

15 The less polar fraction was isolated as a yellow solid (1.2g), a sample of which (815mg) was purified by recrystallisation from toluene (5ml) to give the title compound as colourless needles (600mg). A sample of this material (580mg) was further purified by preparative HPLC using a Luna C8(II) 150x21.2mm 10 μ m column eluting with 95:5 water:acetonitrile (0.1% aqueous trifluoroacetic acid) and acetonitrile (0-1min 100:0 then over 1 min changing to 70:30 for 18min then changing to 100:0 over 1min) to provide a sample of the title compound. This material was freed of any remaining acid by dissolving in dichloromethane (50ml) and washing with saturated aqueous sodium bicarbonate solution (50ml). The organic phase was dried over magnesium sulphate, filtered and concentrated under reduced pressure to provide a foam (270mg) which was recrystallised from toluene (5ml) to give a sample of the title compound as colourless needles (265mg). M.p. 127-128 °C.

^1H NMR (400MHz, CDCl_3): δ = 0.84 (m, 4H), 1.58 (m, 1H), 2.13 (s, 3H), 4.03 (m, 2H), 4.13 (m, 2H), 7.42 (s, 2H), 7.59 (s, 1H).

LRMS (atmospheric pressure chemical ionisation) : m/z [MH^+] 309.


Microanalysis: Found C, 66.14; H, 5.24; N, 18.15. $\text{C}_{17}\text{H}_{16}\text{N}_4\text{O}_2$ requires C, 66.22; H, 5.23; N, 18.17%.

Regioisomer confirmed by nOE NMR and unambiguously identified by X-ray crystallography.



The more polar fraction was further purified by flash chromatography on silica gel eluting with ethyl acetate:toluene (50:50, by volume) to give 5-[[5-cyclopropyl-1-(2-hydroxyethyl)-3-methyl-1*H*-pyrazol-4-yl]oxy]isophthalonitrile (structure below)

5



as a white solid (90mg). M.p. 129-130 °C.

¹H NMR (400MHz, CDCl₃): δ = 0.68 (m, 2H), 0.87 (m, 2H), 1.58 (m, 1H), 2.03 (s, 3H), 4.07 (m, 2H), 4.31 (m, 2H), 7.39 (s, 2H), 7.59 (s, 1H).

10

LRMS (atmospheric pressure chemical ionisation): m/z [MH⁺] 309.

EXAMPLE 2

5-[[3-Cyclopropyl-1-(2-hydroxyethyl)-5-methyl-1*H*-pyrazol-4-yl]oxy]isophthalonitrile


15

To a stirred solution of the pyrazole from Preparation 9 (250mg, 0.64mmol) in methanol (6ml) was added *para*-toluenesulfonic acid (12mg, 0.06mmol). After 18 hours the reaction mixture was concentrated and the residue was partitioned between 10% aqueous potassium carbonate solution (20ml, w/v) and dichloromethane (20ml). The separated aqueous layer was washed with dichloromethane (2 x 20ml) and the combined organic components were dried over magnesium sulfate, filtered and concentrated to give the title compound as a pale yellow oil (195mg) which was used without further purification.

20

¹H NMR (400mHz, CDCl₃) consistent with that described above.

25 LRMS (thermospray): m/z [MH⁺] 309.

PREPARATION 1**1,3-Dibromo-5-methoxybenzene**

5

Sodium methoxide (8.80ml of a 4.5M solution in methanol, 39.6mmol) was added dropwise to a stirred solution of 3,5-dibromofluorobenzene (5.00g, 19.0mmol) (Aldrich) in *N,N*-dimethylformamide (95ml) at 0°C under nitrogen. The reaction was allowed to warm to room temperature, stirred for 1 hour and then concentrated under reduced pressure. The residue was dissolved in ether (500ml) and the resulting solution was washed with water (3x300ml) and brine (300ml), dried over magnesium sulphate, filtered and concentrated under reduced pressure to provide the title compound (5.13g) as a white solid.


15 $^1\text{H-NMR}$ (300MHz, CDCl_3): δ = 3.79 (s, 3H), 7.00 (s, 2H), 7.26 (s, 1H).

LRMS (thermospray): m/z [MH^+] 266.

Microanalysis: Found: C, 31.56; H, 2.29. $\text{C}_7\text{H}_6\text{OBr}_2$ requires C, 31.62; H, 2.27%.

PREPARATION 2

20 **3,5-Dicyanomethoxybenzene**



25 Tris(dibenzylideneacetone)dipalladium(0) (6.53g, 7.15mmol) was added in one portion to a stirred solution of the bromide of Preparation 1 (38.0g, 143mmol) and zinc cyanide (20.0g, 172mmol) in *N,N*-dimethylformamide (300ml) at room temperature under nitrogen. The reaction was heated at 100°C for 14 hours and cooled to room temperature. Water (1500ml) was added and the mixture was extracted with ethyl acetate (3x500ml). The combined organics were filtered and the filtrate was washed with water (500ml), dried over magnesium sulphate, filtered and concentrated under reduced pressure. The resulting solid was

30


trituated with toluene (1000ml) to provide the title compound (18.0g) as a tan solid.

$^1\text{H-NMR}$ (300MHz, CDCl_3): δ = 3.83 (3H, s), 7.31 (2H, s), 7.48 (1H, s).

5

PREPARATION 3

3,5-Dicyanohydroxybenzene



10


The nitrile of Preparation 2 (9.60g, 60.7mmol) was added portionwise to a stirred suspension of aluminium trichloride (32.4g, 243mmol) in dichloromethane (250ml) at 0°C under nitrogen. The suspension was heated to 45°C and stirred for 6 days. The reaction was cooled to room temperature and cautiously poured onto ice (450ml). Concentrated hydrochloric acid (450ml) was added dropwise and the resulting suspension was stirred for 10 minutes at room temperature. The resulting solid was collected by filtration, washed with water and dried over phosphorus pentoxide to provide the title compound (7.83g) as a tan solid containing approximately 11 % starting material by $^1\text{H-NMR}$ and microanalysis.

15

$^1\text{H-NMR}$ (400MHz, CDCl_3): δ = 7.36 (m, 2H), 7.56 (m, 1H).

PREPARATION 4

3-Oxobutanoic acid



25

Sodium hydroxide (37.9g, 0.947mol) was dissolved in water (770ml) and was added dropwise over 20min to 3-oxo-butanoic acid methyl ester (100g, 0.861mol) (Aldrich) at room temperature with stirring. The reaction was stirred for 18 hours, quenched with ammonium sulfate (700g) and acidified slowly with a solution of concentrated hydrochloric acid (21.5ml) in water (250ml) with ice cooling. The reaction mixture was extracted with diethylether (6x200ml) and the combined organic extracts were dried over magnesium sulphate, filtered and concentrated


30

under reduced pressure to provide the title compound (58.2g) as a pale yellow oil which was a mixture of keto:enol tautomers (as observed in ^1H NMR).

^1H NMR (400MHz, CDCl_3): δ = 2.00 (s, 3H-enol), 2.30 (s, 3H-keto), 3.51 (s, 2H-keto), 5.02 (s, 1H-enol).

PREPARATION 5

1-Cyclopropyl-1,3-butanedione




10

Magnesium turnings (3.04g, 125mmol) suspended in methanol (145ml) were heated to reflux under nitrogen for 1 hour, cooled to room temperature and the β -keto acid from Preparation 4 (25.5g, 250mmol) dissolved in methanol (25ml) was added dropwise with ice-cooling. The reaction was stirred for 1 hour at room temperature and the solvent was removed under reduced pressure to give the magnesium salt of the acid. Meanwhile, cyclopropane-carboxylic acid (9.91ml, 125mmol) was dissolved in dimethylformamide (200ml) and carbonyldiimidazole (22.4g, 138mmol) was added portionwise under nitrogen at 0°C . This was stirred for 1.5 hour and the magnesium salt from above was added as a solution in dimethylformamide (100ml) at 0°C . The reaction was allowed to stir at room temperature for 92 hours and the mixture was poured into 2M aqueous hydrochloric acid (85ml) then diluted with water (170ml). The mixture was extracted with diethylether (6x200ml) and the combined organic extracts were washed with brine (3x200ml), dried over magnesium sulphate and concentrated under reduced pressure. The residual orange oil was purified by flash chromatography on silica gel eluting with pentane:diethylether (100:0 changing to 90:10 then 80:20, by volume) to provide the title compound (7.39g) as a yellow oil.

30

^1H NMR (400MHz, CDCl_3): δ = 0.89 (m, 2H), 1.08 (m, 2H), 1.59 (m, 1H), 2.00 (s, 3H), 5.61 (s, 1H), 15.62 (s, 1H).

LRMS (electrospray) : m/z [MNa^+] 149.

PREPARATION 62-Chloro-1-cyclopropyl-1,3-butanedione

5

Chlorotrimethylsilane (18.9ml, 174mmol) was added to a solution of tert-butylammonium bromide (932mg, 2.89mmol) in dry acetonitrile (50ml) under nitrogen at room temperature and the mixture was cooled to 0°C. The diketone from Preparation 5 (7.3g, 57.9mmol) in acetonitrile (36ml) was then added followed by dropwise addition of dry dimethylsulfoxide (12.3ml, 174mmol). The reaction was stirred at 0°C for 1.5 hours and the mixture was diluted with water (500ml), extracted with diethylether (2x200ml and 100ml) and the combined organic extracts were dried over magnesium sulphate, filtered and concentrated under reduced pressure. The residual oil was purified by flash chromatography on silica gel eluting with pentane:diethylether (100:0 changing to 95:5 then 90:10, by volume) to provide the title compound (5.76g) as a colourless oil.


10

15

¹H NMR (400MHz, CDCl₃): δ = 1.04 (m, 2H), 1.18 (m, 2H), 2.27 (s, 3H), 2.42 (m, 1H), 15.78 (s, 1H).

LRMS (electrospray): m/z [M-H⁺] 159.

20

PREPARATION 75-[1-(Cyclopropylcarbonyl)-2-oxopropoxy]isophthalonitrile

25

Cesium carbonate (6.01g, 18.5mmol) and the phenol from Preparation 3 (2.66g, 18.5mmol) were added to a stirred solution of the diketone from Preparation 6 (2.46g, 15.4mmol) in acetone (75ml) under nitrogen at 60°C and the reaction was stirred for 3 hours. After cooling the acetone was removed under reduced

30




pressure and the residue was partitioned between 1N aqueous hydrochloric acid (100ml) and dichloromethane (100ml). The aqueous phase was separated and extracted with dichloromethane (50ml). The combined organic components were dried over magnesium sulphate and concentrated under reduced pressure to give the title compound as a cream solid (4.2g).

^1H NMR (400MHz, CDCl_3): δ = 0.92 (m, 2H), 1.19 (m, 2H), 1.78 (m, 1H), 1.99 (s, 3H), 7.47 (m, 2H), 7.62 (m, 1H).

LRMS (electrospray) : m/z $[\text{M}-\text{H}^+]$ 267.

PREPARATION 8

5-[(3-Cyclopropyl-5-methyl-1H-pyrazol-4-yl)oxy]isophthalonitrile




Hydrazine hydrate (298 μ l, 6.16mmol) was added to a solution of the diketone from Preparation 7 (1.50g, 5.60mmol) in acetic acid (22ml) under nitrogen at room temperature. After stirring at 50 $^{\circ}\text{C}$ for 18 hours, the mixture was allowed to cool to room temperature and concentrated under reduced pressure. The residue was partitioned between 10% aqueous potassium carbonate solution (50ml) and dichloromethane (50ml). The separated aqueous layer was washed twice with dichloromethane (2x50ml). The combined organic components were dried over magnesium sulphate, filtered and concentrated under reduced pressure to provide a pale yellow oil. The crude product mixture was purified by flash chromatography on silica gel eluting with pentane:ethyl acetate (75:25 changing to 67:33, by volume) to provide the title compound (1.20g) as a pale yellow oil.

^1H NMR (400MHz, CDCl_3): δ = 0.75 (m, 2H), 0.85 (m, 2H), 1.60 (m, 1H), 2.10 (s, 3H), 7.40 (s, 2H), 7.60 (s, 1H).

LRMS (thermospray): m/z $[\text{MH}^+]$ 260.

PREPARATION 9

5-({3-Cyclopropyl-5-methyl-1-[2-(tetrahydro-2H-pyran-2-yloxy)ethyl]-1H-pyrazol-4-yl}oxy)isophthalonitrile



5

To a stirred solution of the pyrazole from Preparation 8 (420mg, 1.59mmol) in dimethylformamide (4ml) at 0°C was added sodium hydride (70mg of a 60% w/w suspension in mineral oil). After the addition was complete the cooling bath was removed and the mixture was stirred at room temperature for 30 minutes. A solution of 2-(2-bromoethoxy)tetrahydro-2H-pyran (264μl, 1.75mmol) in dimethylformamide (2ml) was added. After 2 hours the reaction mixture was quenched by addition of water (20ml) and was extracted with dichloromethane (3 x 20ml). The combined organic components were washed with brine (2 x 20ml), dried over magnesium sulfate, filtered and concentrated to give a yellow oil. The crude product mixture was purified by flash chromatography on silica gel eluting with dichloromethane:methanol (100:0 changing to 98:2, by volume) to provide a mixture of the two regioisomers (594mg). The two regioisomers were separated by flash chromatography on silica gel eluting with toluene:ethyl acetate (100:0 changing to 80:20, 75:25, 67:33 and 50:50 by volume) to provide the title compound (257mg) (less polar fraction) and its regioisomer (90mg) (more polar fraction).

Less Polar Fraction


¹H NMR (400MHz, CDCl₃): δ = 0.78 (m, 4H), 1.55 (m, 5H), 1.67 (m, 2H), 2.12 (s, 3H), 3.45 (m, 1H), 3.65 (m, 1H), 3.75 (m, 1H), 4.04 (m, 1H), 4.18 (m, 2H), 4.53 (m, 1H), 7.40 (s, 2H), 7.59 (s, 1H).

LRMS (thermospray): m/z [MH⁺] 393.



More Polar Fraction

5-({5-Cyclopropyl-3-methyl-1-[2-(tetrahydro-2H-pyran-2-yloxy)ethyl]-1H-pyrazol-4-yl}oxy)isophthalonitrile




- 5 ¹H NMR (300MHz, CDCl₃): δ = 0.68 (m, 2H), 0.85 (m, 2H), 1.53 (m, 3H), 1.72 (m, 4H), 2.10 (s, 3H), 3.51 (m, 1H), 3.72 (m, 1H), 3.83 (m, 1H), 4.17 (m, 1H), 4.35 (m, 2H), 4.58 (m, 1H), 7.38 (s, 2H), 7.59 (s, 1H).
LRMS (thermospray): m/z [MH⁺] 393.

09/08/20



CLAIMS

1. The compound of formula (I)



5

or a pharmaceutically acceptable salt, solvate or derivative thereof.

2. A pharmaceutical composition including the compound of formula (I) or a pharmaceutically acceptable salt, solvate or derivative thereof together with one or more pharmaceutically acceptable excipients, diluents or carriers.

10

3. A pharmaceutical composition according to claim 2 including one or more additional therapeutic agents.

4. The compound of formula (I) or a pharmaceutically acceptable salt, solvate or derivative thereof, or a pharmaceutical composition according to claim 2 or 3, for use as a medicament.

15

5. The compound of formula (I) or a pharmaceutically acceptable salt, solvate or derivative thereof, or a pharmaceutical composition according to claim 2 or 3, for use as a reverse transcriptase inhibitor or modulator.

20

6. The compound of formula (I) or a pharmaceutically acceptable salt, solvate or derivative thereof, or a pharmaceutical composition according to claim 2 or 3, for use in the treatment of an HIV or genetically-related retroviral infection, or a resulting acquired immune deficiency syndrome (AIDS).

25

7. Use of the compound of formula (I) or a pharmaceutically acceptable salt, solvate or derivative thereof, or a pharmaceutical composition according to claim 2 or 3, for the manufacture of a medicament having reverse transcriptase inhibitory or modulating activity.

30




8. Use of the compound of formula (I) or a pharmaceutically acceptable salt, solvate or derivative thereof, or a pharmaceutical composition according to claim 2 or 3, for the manufacture of a medicament for the treatment of an HIV or genetically-related retroviral infection, or a resulting acquired immune deficiency syndrome (AIDS).

9. A method of treating an HIV or a genetically-related retroviral infection, or a resulting acquired immune deficiency syndrome (AIDS), comprising administering an effective amount of the compound of formula (I) or a pharmaceutically acceptable salt, solvate or derivative thereof, or a pharmaceutical composition according to claim 2 or 3.

10. A process for preparing the compound of formula (I) or a salt, solvate or pharmaceutically acceptable derivative thereof, which comprises:


15

(A) condensation of a compound of formulae (II), (VI) or (VII)



with the compound of formula (V)


20



or a salt or hydrate thereof;




(B) alkylation of a pyrazole of formula (XIII)



with an alkylating agent of formula (XIV)

5



or a protected derivative thereof;

10 (C) deprotecting a protected derivative of the compound of formula (I);

and optionally converting the compound of formula (I) prepared by any one of processes (A) to (C) into pharmaceutically acceptable salt, solvate or derivative thereof.


15

11. A compound of formulae (II), (VI), (VII), (VIII), (IX) or (XIII).



ABSTRACT

This invention relates to the pyrazole derivatives of formula (I)



and pharmaceutically acceptable salt, solvate or derivatives thereof, to their use in medicine, to compositions containing them, to processes for their preparation and to intermediates used in such processes.

The compounds of the invention bind to the enzyme reverse transcriptase and are modulators, especially inhibitors, thereof. Reverse transcriptase is implicated in the infectious lifecycle of Human Immunodeficiency Virus (HIV). Compounds which interfere with the function of this enzyme have shown utility in the treatment of conditions caused by HIV and genetically related retroviruses, such as Acquired Immune Deficiency Syndrome (AIDS).

